

**Supplementary for ICD-10 codes used for our analysis**  
**“Characteristics and clinical outcomes of patients with Sarcoidosis admitted for ST-elevation myocardial infarction.”**

Sarcoidosis	D86x
STEMI	I2101, I2102, I2109, I2111, I2119, I2121, I2129, I213
Diabetes	E08.X-E13.X
Hypertension	I10
Smoking	F17-F17299, Z720, Z87891, Z716
CKD	N18x
Dyslipidemia	E780 E7800 E7801 E781 E782 E783 E784 E7841 E7849 E785
Cirrhosis	K740 K7400 K7401 K7402 K741 K742 K743 K744 K745 K7460 K7469 K7030 K7031 P7881 K717
Prior stroke	Z8673
PVD	I70.X, I73.9
Old MI	I252
Alcohol abuse	Based on Elixhauser
Drug abuse	Based on Elixhauser
AIS	I63x
Cardiogenic shock	R570
AKI	N17x
Depression	Based on Elixhauser
CABG	Based on CCS CAR003
PCI	Based on CCS CAR004
Atrial fibrillation	I480, I481, I4811, I4819, I482, I4820, I4821, I4891
SVT	I470, I471, I479
VT	I472
Ventricular fibrillation	I4901, I462, I468, I469
Atrial flutter	I483, I484, I4892